



**INSTRUCTIONS FOR COLONOSCOPY WITH SUTAB  
SPLIT – DOSE**

**MEDICATIONS:** Please notify us if you are on the following medications: **Aggrenox, Plavix, Coumadin, Warfarin, Pradaxa, Effient, Xarelto or Eliquis.** You may need to stop these medications 5 days prior to your procedure. Lovenox or Fragmin should not be used the day of the procedure. If you are diabetic, please consult your primary care physician regarding your medication and diet. You may continue to take all other medications. Please call our office with any questions that you may have.

**Commercial Sutab coupon to show pharmacy:**

**Rx. BIN: 004682**

**PCN: CN**

**GROUP: WCSEB4105**

**MEMBER ID: 42173175453**

**Medicare link for coupon:**

**Patients can access and activate it online for coupon**

**<https://sutab.com/Savings>**

**LOW FIBER DIET 3 DAYS PRIOR TO COLONOSCOPY- PLEASE SEE BELOW AT BOTTOM**

**LIQUID DIET:** The day before colonoscopy, consume CLEAR LIQUIDS ONLY! DO NOT EAT ANYTHING! Avoid dairy products and anything that is red, purple or blue!! Drink plenty of fluids all day.

**RESTRICTED LIQUID DIET INCLUDES: (NO RED, BLUE, OR PURPLE PLEASE!)**

Strained, pulp free fruit juices (apple, white grape, lemonade)	Water
Coffee or tea (no milk or creamer)	Gatorade
Carbonated/non-carbonated soft drinks	Popsicles
Kool-Aid or other fruit flavored drinks	Clear broth or bouillon
Plain Jell-O (no added toppings or fruit)	NO dairy products or alcohol

**PREP INSTRUCTIONS: PLEASE CONTACT YOUR GASTROENTEROLOGIST FOR SPECIFIC DOSE TIMES**

PREP INSTRUCTIONS:

**SUTAB** is a split-dose regimen. A total of 24 tablets is required for complete preparation for colonoscopy. You will take the tablets in two doses of 12 tablets each. Water must be consumed with each dose of SUTAB, and additional water must be consumed after each dose.

**DOSE 1** - On the Day Prior to Colonoscopy

**STEP 1** Open 1 bottle of 12 tablets. Take the tablets at **(Please contact GI MD for exact time)** with water



**STEP 2** Fill the provided container with **16 ounces of water** (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15 to 20 minutes. Drink additional water.

**STEP 3** Approximately **1 hour after** the last tablet is ingested, fill the provided container again with **16 ounces of water** (up to the fill line), and drink the entire amount **over 30 minutes**.

**STEP 4** Approximately **30 minutes** after finishing the second container of water, fill the provided container with **16 ounces of water** (up to the fill line), and drink the entire amount **over 30 minutes**.

**DAY PRIOR TO PROCEDURE @ (Please contact GI MD for exact time):** REPEAT Step 1 to Step 4

**YOU SHOULD NOT HAVE ANYTHING TO EAT OR DRINK 4 HOURS PRIOR TO THE PROCEDURE.**

**TRANSPORTATION:** Taxicabs and UBER ARE NOT acceptable. You must have someone you know drive you home. From the time you arrive at our facility until the time you leave, may require 2 to 3 hours.

**CALL THE OFFICE 48 HOURS IN ADVANCE IF YOU NEED TO CANCEL OR RESCHEDULE THE PROCEDURE. 469-800-6020**

**FAILURE TO CANCEL OR RESCHEDULE PROCEDURES WITHIN 48 HOURS WILL RESULT IN A \$100 NO SHOW FEE**

**IMPORTANT BILLING INFORMATION:**

We will bill your health insurance carrier if you have provided us with complete information at the time of registration. However, please remember you are ultimately responsible for payments at Time of Service.

**LOW FIBER DIET 3 DAYS PRIOR TO COLONOSCOPY- PLEASE SEE BELOW AT BOTTOM**

## Low-Residue/Low-Fiber Diet

A low-residue/low-fiber diet limits the amount of food waste that has to move through the large intestine.

### Points to Keep in Mind

- Avoid any food made with seeds, nuts, or raw or dried fruit.
- Avoid whole-grain breads and cereals. Purchase products made from refined flour.
- Do not eat raw fruits or vegetables. Remove skins before cooking.

Types of Foods	Foods to Choose	Foods to Avoid
Breads, Cereals, Rice, and Pasta	<ul style="list-style-type: none"> <li>▪ Enriched white bread, rolls, biscuits, and muffins</li> <li>▪ Waffles, French toast, and pancakes</li> </ul>	<ul style="list-style-type: none"> <li>▪ Breads or rolls with nuts, seeds, or fruit</li> </ul>

	<ul style="list-style-type: none"> <li>White rice, noodles, pasta, and cooked potatoes (no skin)</li> <li>Plain crackers</li> <li>Cooked cereals: farina, cream of wheat, and grits</li> <li>Cold cereals: Puffed Rice, Rice Krispies, Corn Flakes, and Special K</li> </ul>	<ul style="list-style-type: none"> <li>Whole wheat, pumpernickel bread, and cornbread</li> <li>Potatoes with skin, brown or wild rice, and kasha (buckwheat)</li> <li>Whole grain cereals, bran cereals, granola-type cereals, and cereals with nuts, seeds, coconut, or dried fruit</li> </ul>
<b>Fruit</b>	<ul style="list-style-type: none"> <li>Strained fruit juice</li> <li>Canned or cooked fruits without skins or seeds</li> <li>Ripe banana</li> <li>Soft cantaloupe and honeydew melon</li> </ul>	<ul style="list-style-type: none"> <li>Prunes and prune juice</li> <li>Raw or dried fruit</li> <li>All berries and raisins (especially raspberries and blackberries)</li> </ul>
<b>Milk and Dairy Products</b>	<ul style="list-style-type: none"> <li>Milk, plain or flavored</li> <li>Yogurt, custard, and ice cream</li> <li>Cheese and cottage cheese</li> </ul>	<ul style="list-style-type: none"> <li>Yogurt with nuts or seeds</li> </ul>
<b>Vegetables</b>	<ul style="list-style-type: none"> <li>Strained vegetable juice</li> <li>Well-cooked fresh or canned vegetables such as asparagus tips, beets, green beans, carrots, acorn squash (without seeds), pureed spinach, and tomato sauce</li> </ul>	<ul style="list-style-type: none"> <li>Raw or partially cooked (steamed) vegetables</li> <li>Vegetables with seeds</li> <li>Sauerkraut</li> <li>Cooked peas, winter squash, broccoli, brussels sprouts, cabbage, onions, cauliflower, baked beans and corn</li> </ul>
<b>Meats, Poultry, Fish, Dry Beans, Peas, and Eggs</b>	<ul style="list-style-type: none"> <li>Ground, well-cooked, tender beef, lamb, ham, veal, pork, fish, poultry, and organ meats</li> <li>Eggs</li> </ul>	<ul style="list-style-type: none"> <li>Dry beans, peas, and lentils</li> <li>Peanut butter</li> </ul>
<b>Fats, Snacks, Sweets, Condiments and Beverages</b>	<ul style="list-style-type: none"> <li>Margarine, butter, oils, mayonnaise, sour cream, and salad dressing</li> <li>Plain gravies</li> <li>Sugar, clear jelly, honey, and syrup</li> <li>Spices, cooked herbs, bouillon, broth, and soups made with allowed ingredients</li> <li>Coffee, tea, and carbonated drinks</li> <li>Plain cakes and cookies</li> </ul>	<ul style="list-style-type: none"> <li>Nuts, seeds, and coconut</li> <li>Jam, marmalade, and preserves</li> <li>Pickles, olives, relish, and horseradish</li> <li>All desserts containing nuts, seeds, dried fruit, coconut, or made from whole grains or bran</li> <li>Candy made with nuts or seeds</li> </ul>



- Gelatin, plain puddings, custard, ice cream, ▪ Popcorn  
sherbet, Popsicles
  - Hard candy
  - Pretzels
-